arXiv:math/0306368v2 [math.DG] 29 Jun 2003

KÄHLER SOLVMANIFOLDS

DONU ARAPURA

Abstract. Compact Kähler solvmanifolds are classified up to biholomorphism. A proof of a conjecture Benson and Gordon, that completely solvable compact Kähler solvmanifolds are tori is deduced from this.

In this note, we use the term solvmanifold in a somewhat restricted sense. A solvmanifold will be a compact quotient G/Γ of a connected simply connected solvable real Lie group by a discrete subgroup. Such a space is a $K(\Gamma, 1)$. Our goal is to classify all Kähler solvmanifolds up to biholomorphism. To be precise, we consider a Kähler solvmanifold to be a solvmanifold with complex structure and Kähler metric that are not assumed invariant under the group. (The invariance of the complex structure will follow from our theorem.) Complex tori are clearly examples of such spaces. Hasegawa [H2] has shown how to construct additional examples by taking quotients of suitable tori, and we show that all Kähler solvmanifolds arise this way. As a corollary, we prove a conjecture of Benson and Gordon [BG2] that tori are the only Kähler solvmanifolds corresponding to completely solvable Lie groups[1]. The main new ingredient here is the restriction theorem for polycyclic Kähler groups proved by Nori and the author [AN].

After the completion of the first version of this paper, I have learned from Alex Brudnyi that he had obtained similar results in his 1995 PhD thesis (Technion) and subsesquent papers [Br1, Br2]. However the methods are quite different and do not yield the analogous result for solvmanifolds which are bimeromorphic to a compact Kähler manifolds (theorem 2).

Let us start by defining a few basic terms and constructions. A solvable group G is called completely solvable, or of real type, if all the eigenvalues of images of the elements of its Lie algebra under its adjoint representation are real. Given a solvmanifold G/Γ, the map

$$G/\Gamma \to \frac{G/[G,G]}{im(\Gamma \to G/[G,G])}$$

defines a fiber bundle over a torus called the canonical torus fibration in [AS].

Lemma 1. *The fibers are compact nilmanifolds (i.e. quotients of connected nilpotent Lie groups by lattices).*

Proof. Since $[G,G]$ is nilpotent [R, 3.11], it lies in the maximum connected nilpotent subgroup N. $N \cap \Gamma$ is a cocompact lattice in N by a theorem of Mostow [M], see also [R, 3.3]. Hence $[G,G] \cap \Gamma$ is also a cocompact lattice in $[G,G]$ [R, 2.3]. □

A complex torus is a quotient V/L of a complex vector space by a lattice. We can identify V with the holomorphic tangent space at 0, and the map $V \to V/L$

Partially supported by the NSF.

[1] The proof in [TK] is incomplete; see [H2, p.257].

with the exponential map. Holomorphic maps of complex tori can always be lifted to affine maps of vector spaces [GH, p.326]; in particular, maps preserving 0 are necessarily homomorphisms. For any compact Kähler manifold X, the Albanese torus $Alb(X)$ is a complex torus with real dimension equal to the first Betti number of X [GH, p. 331]. After choosing a base point $x \in X$, there is a holomorphic map, called the Abel-Jacobi map, $\alpha : X \to Alb(X)$ with $\alpha(x) = 0$ which is topologically just the classifying map $X \to K(H_1(X, \mathbb{Z})/torsion, 1)$.

Lemma 2. *If a compact Kähler manifold is diffeomorphic to a torus then it is biholomorphic to a complex torus.*

Proof. Let X be the manifold in question. Then it follows that the Albanese map is a holomorphic diffeomorphism and hence biholomorphism. □

When X is a Kähler solvmanifold, the Abel-Jacobi map would coincide with the canonical torus fibration. We will make use of this fact later on.

We give a general method for producing Kähler solvmanifolds along the lines of Hasegawa's paper [H2]. This method will produce all examples. We will make use of the standard description of group extensions using cohomology [B].

Construction 1. *Let*

$$0 \to \mathbb{Z}^{2m} \to \Gamma \to \mathbb{Z}^{2n} \to 0$$

be a group extension such that

(a) *The homomorphism $\mathbb{Z}^{2n} \to GL(2m, \mathbb{Z})$, associated to the extension, has finite image.*
(b) *$\mathbb{Z}^{2n} \to GL(2m, \mathbb{Z})$ extends to a homomorphism $\mathbb{R}^{2n} \to GL(2m, \mathbb{R})$ of real Lie groups.*
(c) *There is a complex structure J_0 on $\mathbb{R}^{2m}$ invariant under the action of $\mathbb{Z}^{2n}$.*
(d) *The class of the extension in $H^2(\mathbb{Z}^{2n}, \mathbb{Z}^{2m})$ has finite order.*

Let $V = \mathbb{R}^{2m} \oplus \mathbb{R}^{2n}$, and choose complex structure J on it extending J_0. Let $\Gamma \otimes \mathbb{R}$ denote the semidirect product $\mathbb{R}^{2m} \rtimes \mathbb{R}^{2n}$ under the action (b). Note that as a sets, $\Gamma \otimes \mathbb{R}$ and V are one and the same, nevertheless it is useful to keep a distinction. Consider the diagram

$$\begin{tikzcd}
0 \arrow[r] & \mathbb{Z}^{2m} \arrow[r] \arrow[d] & \Gamma \arrow[r] \arrow[d] & \mathbb{Z}^{2n} \arrow[r] \arrow[d] & 0 \\
0 \arrow[r] & \mathbb{R}^{2m} \arrow[r] & \tilde{\Gamma} \arrow[r] & \mathbb{Z}^{2n} \arrow[r] & 0
\end{tikzcd}$$

where the left hand square is the "pushout". Assumption (d) shows that the bottom sequence splits. Therefore we get embeddings $\Gamma \subset \tilde{\Gamma} \subset \Gamma \otimes \mathbb{R}$. The quotient $X = (\Gamma \otimes \mathbb{R})/\Gamma$ is a solvmanifold. Under the identification $\Gamma \otimes \mathbb{R} = V$, Γ will act on V by complex affine transformations. Therefore X inherits a complex structure. We can find, thanks to (a), a Γ-invariant Hermitean inner product on V, and this descends to a flat Kähler metric on X. Note that different choices of J can yield nonbiholomorphic, but diffeomorphic, examples.

Examples of the above type can be constructed quite explicitly. For instance, they include all the hyperelliptic (also called bielliptic) surfaces [H2]. In particular, they not all tori. However, they are always quotients of tori by a finite group acting freely. This is a consequence of the next lemma which implies that Γ contains a normal abelian subgroup Δ of finite index. Then X is a quotient of the complex torus V/Δ by Γ/Δ.

Lemma 3. *Suppose that* Γ *is a finitely generated group which fits into an extension,*

$$0 \to \mathbb{Z}^{2m} \to \Gamma \xrightarrow{\pi} \mathbb{Z}^{2n} \to 0$$

Then conditions (a) and (d), above, hold if and only if Γ *contains a normal abelian subgroup of finite index.*

Proof. Suppose conditions (a) and (d) hold. Let Δ be the preimage in Γ of $\Delta' = ker[\mathbb{Z}^{2n} \to GL(2m, \mathbb{Z})]$. This has finite index. Since $H^2(\Delta', \mathbb{Z}^{2m})$ is torsion free [B], (d) shows that the sequence

$$0 \to \mathbb{Z}^{2m} \to \Delta \to \Delta' \to 0$$

splits. Therefore $\Delta \subset \Gamma$ is a normal abelian subgroup of finite index.

Conversely, let $\Delta \subset \Gamma$ be a normal abelian subgroup of finite index. Since $\pi(\Delta)$ acts trivially on $\Delta \cap \mathbb{Z}^{2m}$ which has finite index in $\mathbb{Z}^{2m}$, it follows that it acts trivially on $\mathbb{Z}^{2m}$. Consequently $\mathbb{Z}^{2n} \to GL(m, \mathbb{Z})$ factors through $\mathbb{Z}^{2n}/\pi(\Delta)$. Let $\tilde{\Delta} = \pi^{-1}(\pi(\Delta))$ and consider the commutative diagram

$$\begin{tikzcd}
0 \arrow[r] & \mathbb{Z}^{2m} \arrow[r] & \Gamma \arrow[r] & \mathbb{Z}^{2n} \arrow[r] & 0 \\
0 \arrow[r] & \mathbb{Z}^{2m} \arrow[r] \arrow[u, equal] & \tilde{\Delta} \arrow[r] \arrow[u] & \pi(\Delta) \arrow[r] \arrow[u] & 0 \\
0 \arrow[r] & \Delta \cap \mathbb{Z}^{2m} \arrow[r] \arrow[u] & \Delta \arrow[r] \arrow[u] & \pi(\Delta) \arrow[r] \arrow[u, equal] & 0
\end{tikzcd}$$

Since Δ is abelian, the extension class of the bottom row is 0. The lower left hand square is a pushout. Therefore the extension class of the bottom row maps to the class of the middle row, and consequently this is also zero. The upper right hand square is a pullback. This implies that the extension class e of the top row maps to the class of the middle row under the map $H^2(\mathbb{Z}^{2n}, \mathbb{Z}^{2m}) \to H^2(\pi(\Delta), \mathbb{Z}^{2m})$. This homomorphism is injective modulo torsion, since the transfer map [B] gives a splitting after tensoring with $\mathbb{Q}$. Therefore e must be torsion. $\square$

Theorem 1. *Every Kähler solvmanifold is biholomorphic to one of the examples described in Construction 1.*

Remark 1. *The theorem is a classification up to biholomorphism not isometry. These examples can have nonflat Kähler metrics, and there does not appear to be any sensible way to classify these.*

Proof of theorem. Let Γ be a cocompact lattice in a simply connected solvable Lie group G such that $X = G/\Gamma$ has a Kähler structure. By [R, 4.28], Γ is polycyclic. Therefore [AN, 4.10] implies that Γ contains a nilpotent subgroup $\Delta_1 \subseteq \Gamma$ of finite index. We can find a subgroup $\Delta \subseteq \Delta_1$ of finite index which is normal in Γ. Δ is again nilpotent, thus $T = \Gamma/\Delta$ is an aspherical manifold with a nilpotent fundamental group. Pulling back the Kähler structure from X also makes T into a Kähler manifold such that the projection $T \to X$ is holomorphic. Therefore by [BG1, theorem B], T is homotopic to a torus. Consequently, [M, theorem A] shows that T is diffeomorphic to a torus. Then T is a complex torus by lemma 2.

Consider the Abel-Jacobi map $\alpha : X \to A = Alb(X)$. This is surjective since it coincides with the canonical torus fibration. Let $f : T \to A$ denote the composition $T \to X \to A$. We can choose a base point of X so that $0 \in T$ maps to $0 \in A$. Then $f : T \to A$ becomes a homomorphism. Therefore f is a bundle with fiber $K = ker(f)$ which is a product of a torus with a finite abelian group. In particular,

$T \to A$ is a submersion, which implies that $X \to A$ is also a submersion. Therefore the fibers of α are connected Kähler nilmanifolds (lemma 1), and hence tori by [BG1, theorem A] or [H1]. It follows also that $[G, G]$ is abelian, and hence isomorphic to a Euclidean space (with even dimension).

The exact sequence for homotopy groups of a fibration applied to α yields the extension

$$0 \to \mathbb{Z}^{2m} \to \Gamma \to \mathbb{Z}^{2n} \to 0$$

We have to check that the conditions of construction 1 hold. Properties (a) and (d) follow from lemma 3. The action of of $G/[G, G] \cong \mathbb{R}^{2n}$ on $[G, G] \cong \mathbb{R}^{2m}$ gives the extension required for (b). Let $V = \mathbb{R}^{2m} \oplus \mathbb{R}^{2n}$, Γ acts on this by real affine transformations. V has a complex structure J coming from the identification of V with the tangent space of T at 0. $\mathbb{R}^{2m}$ is invariant under J since it can be identified with the tangent space to K at 0. The action of Γ on V is holomorphic with respect to J, therefore Γ acts by complex affine transformations. In particular, the action of $\mathbb{Z}^{2n}$ on $\mathbb{R}^{2m}$ leaves $J|_{\mathbb{R}^{2m}}$ invariant, and this completes the proof. □

As corollaries, we confirm conjectures of Hasegawa and Benson and Gordon.

Corollary 1. *A Kähler solvmanifold is both a quotient of a complex torus and a bundle of complex tori over a complex torus.*

Proof. In the notation of the previous proof, X is quotient of T by Γ/Δ. We have a submersion $\alpha : X \to A$. The map $T \to X$ induces a surjective homomorphism from K to a fiber of α with a finite kernel H. From this, we can deduce that $X \to A$ is a locally trivial bundle with fiber K/H. □

Corollary 2. *Let $X = G/\Gamma$ be a Kähler solvmanifold with G completely solvable. Then X is biholomorphic to a complex torus.*

Proof. The theorem shows that G contains a lattice which is free abelian of even rank. At this point, we are in a position to argue as in [TK]. We can apply Saito's rigidity theorem [S] to conclude that G is in fact abelian, and therefore that X is a torus. □

We can prove a slightly stronger result by the same method.

Theorem 2. *The examples given in Construction 1 are the only complex solvmanifolds which are bimeromorphic to a compact Kähler manifolds.*

It is just a matter of going through each step and making sure that everything still works. We indicate the main points:

- The key step that a nilmanifold cannot be bimeromorphic to a compact Kähler manifolds unless it is a torus is already supplied by Hasegawa [H1]. This is deduced by showing that a nilmanifold cannot be formal unless it is a torus and then applying [DGMS]. This argument yields a bit more, namely a generalization of [BG1, theorem B], that tori are the only compact bimeromorphically Kähler manifolds homotopic to a nilmanifold.
- The above discussion about the Albanese carries over to the class compact bimeromorphically Kähler manifolds without any modification. See for example [GPR, p. 337]

- Since fundamental groups of complex manifolds are invariant under blow ups with smooth center, the classes of fundamental groups of compact bimeromorphically Kähler manifolds and compact Kähler manifolds coincide. Thus [AN, 4.10] also applies to the former.

References

[AN] D. Arapura, M. Nori *Solvable fundamental groups of algebraic varieties and Kähler manifolds*, Compositio Math. (1999)

[AS] L. Auslander, M. Auslander, *Solvable Lie groups and locally Euclidean Riemannian spaces*, Proc. AMS (1958)

[AS] L. Auslander, R. Szczarba *Vector bundles over tori and noncompact solvmanifolds*, Amer. J. Math (1975)

[BG1] C. Benson, C. Gordon, *Kähler and symplectic nilmanifolds*, Topology (1988)

[BG2] C. Benson, C. Gordon, *Kähler structures on compact solvmanifolds*, Proc. AMS (1990)

[B] K. Brown, *Cohomology of groups*, Springer-Verlag (1982)

[Br1] A. Brudnyi, *Classification theorem for a class of flat connections and representations of Khler groups*, Michigan Math. J. 46 (1999)

[Br2] A. Brudnyi, *Solvable representations of Kähler groups*, preprint arXiv.math.CV/9911080

[DGMS] P. Deligne, P. Griffiths, J. Morgan, D. Sullivan, *Real homotopy theory of Kähler manifolds*, Invent. Math. 29 (1975)

[GH] P. Griffiths, J. Harris *Principles of algebraic geometry*, Wiley (1978)

[GPR] H. Grauert, T. Peternell, R. Remmert (eds.), *Several Complex Variables VI*, Springer-Verlag (1991)

[H1] K. Hasegawa *Minimal models of nilmanifolds*, Proc. AMS (1989)

[H2] K. Hasegawa *A class of compact Kählerian solvmanifolds and a general conjecture*, Geom. Ded. (1999)

[M] G. Mostow *Factor spaces of solvable Lie groups*, Ann. Math (1954)

[R] M. S. Raghunathan, *Discrete subgroups of Lie groups*, Springer-Verlag (1972)

[S] M. Saito, *Sur certains groupes de Lie résolubles. II*, Sci. Papers Coll. Gen. Ed. Univ. Tokyo 7 (1957)

[TK] A. Tralle, W. Kedra, *Compact completely solvable Kähler solvmanifolds are tori*, Internat. Math. Res. Not. (1997)

Department of Mathematics, Purdue University, West Lafayette, IN 47907, U.S.A.
E-mail address: `arapura@math.purdue.edu`